



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

288. Proposed by DR. L. E. DICKSON, Associate Professor of Mathematics, The University of Chicago.

Evaluate the determinant which arises in finding the inverse of the transformation, with binomial coefficients,

$$T: \quad \xi_i = \sum_{j=i}^{g-1} \binom{j}{i} x_j \quad (i=0, 1, \dots, g-1).$$

Solution by the PROPOSER.

Denote by  $D_{n,m}$  the minor of the element in the  $(n+1)$ th row and  $(m+1)$ th column. Evidently  $D_{nn}=1$ ,  $D_{nm}=0$  for  $n < m$ . For  $n=m+k$ ,  $k > 0$ ,

$$D_{m+k, m} = \begin{vmatrix} \binom{m+1}{1} & \binom{m+2}{2} & \binom{m+3}{3} \dots \binom{m+k}{k} \\ 1 & \binom{m+2}{1} & \binom{m+3}{2} \dots \binom{m+k}{k-1} \\ 0 & 1 & \binom{m+3}{1} \dots \binom{m+k}{k-2} \\ \cdot & \cdot & \cdot \dots \cdot \\ 0 & 0 & 0 \dots \binom{m+k}{1} \end{vmatrix}$$

Studnicka\* has evaluated a similar determinant:

$$\begin{vmatrix} \binom{m+1}{1} & \binom{m+1}{2} & \binom{m+1}{3} \dots \binom{m+1}{k} \\ 1 & \binom{m+1}{1} & \binom{m+1}{2} \dots \binom{m+1}{k-1} \\ \cdot & \cdot & \cdot \dots \cdot \\ 0 & 0 & 0 \dots \binom{m+1}{1} \end{vmatrix} = \binom{m+k}{k}.$$

Multiplying the first  $j-1$  columns of the latter by, respectively,

$$\binom{j-1}{j-1}, \binom{j-1}{j-2}, \binom{j-1}{j-3}, \dots, \binom{j-1}{1},$$

and add the products to the  $j$ th column. Then the  $i$ th element in the new  $j$ th column is† (for  $s=j-r$ ),

$$\sum_{r=i-1}^j \binom{m+1}{r-i+1} \binom{j-1}{j-r} = \sum_{s=0}^{j-i+1} \binom{m+1}{j-i+1-s} \binom{j-1}{s} = \binom{m+j}{j-1+1}.$$

Taking  $j=k$ ,  $k-1$ , ..., 2, in turn, we obtain  $D_{m+k, m}$ . Hence

\*Cf. Pascal-Leitzmann, *Die Determinanten*, 1900, p. 134.

†Cf. Netto, *Combinatorik*, p. 250, (19); Hagen, *Synopsis*, Vol. 1, p. 65, 5.

$$D_{m+k, m} = \binom{m+k}{k}, \quad D_{nm} = \binom{n}{m} \text{ if } n > m.$$

Hence the inverse of transformation  $T$  is

$$T^{-1} : x_i = \sum_{j=i}^{g-1} (-1)^{i+j} \binom{j}{i} \xi_j \quad (i=0, 1, \dots, g-1).$$

Since the product of the two transformations is the identity, we have

$$\sum_{j=i}^l (-1)^{i+j} \binom{l}{j} \binom{j}{i} = \delta_{il} \quad (\delta_{ii}=1, \delta_{ii}=0 \text{ if } i \neq l).$$

Conversely, from this well known formula (cf. Netto, p. 255, (43)), follows the evaluation of the determinant  $D$ .

289. Proposed by S. A. COREY, Hiteman, Iowa.

Prove that  $\left( \frac{n+1}{(n+1)^2-1} + \frac{1}{3} \cdot \frac{n+3}{(n+3)^2-1} + \frac{1}{5} \cdot \frac{n+5}{(n+5)^2-1} + \dots \right)$   
 $+ \left( \frac{2}{3} \cdot \frac{1}{n+2} + \frac{4}{15} \cdot \frac{1}{n+4} + \frac{8}{35} \cdot \frac{1}{n+6} + \dots \right) = \frac{n-1}{(n-1)^2-1} + \frac{1}{3} \cdot \frac{n-3}{(n-3)^2-1}$   
 $+ \frac{1}{5} \cdot \frac{n-5}{(n-5)^2-1} + \dots + \frac{1}{l} \cdot \frac{n-l}{(n-l)^2-1},$   $n$  being any odd integer greater than 1  
 and  $l=n-2$ .

Solution by the PROPOSER.

The term by term product of the right hand members of the two Fourier's sine series,

$$1 = \frac{4}{\pi} \left[ \frac{\sin x}{1} + \frac{\sin 3x}{3} + \frac{\sin 5x}{5} + \dots \right] \text{ and}$$

$$\cos x = \frac{4}{\pi} \left[ \frac{2}{3} \cdot \sin 2x + \frac{4}{15} \cdot \sin 4x + \frac{8}{35} \cdot \sin 6x + \dots \right]$$

being equal to the product of the left hand members, may be thus written:

$$\cos x = C_0 + C_1 \cos x + C_2 \cos 2x + C_3 \cos 3x + \dots$$

the  $C$ 's being constants and functions of the coefficients of the sine terms. But as this product of the right hand members is of the same form as the regular Fourier's cosine series for  $\cos x$ , it must be identical with the latter series, and, hence, each  $C$  except  $C_1$  must be equal to zero. Each  $C$  with